

## FOR ANIMAL TREATMENT ONLY

# EIMERIAVAX™ 3m

**Active Constituents:** This vaccine contains viable oocysts of *Eimeria acervulina* Strain RA<sub>3+20</sub>, *E. maxima* Strain MCK<sub>+10</sub> and *E. tenella* Strain Rt<sub>3+15</sub> suspended in phosphate buffered saline (PBS). Each dose comprises a minimum of *E. acervulina* 50 oocysts, *E. maxima* 100 oocysts and *E. tenella* 150 oocysts, with a minimum predicted titre of  $1.2 \times 10^4$  oocysts per mL for 1000 dose and  $6.0 \times 10^4$  oocysts per mL for 5000 dose at the end of the shelf-life.

**Statement of Claims:** Live oocyst preparation to aid in the control of three of the species of *Eimeria* that cause Coccidiosis in breeders, broilers and layer chickens in Australia.

## 1000 / 5000 Doses

**Contents:** 1000 or 5000 doses in a plastic bottle. The *Eimeria* oocysts are supplied as a suspension of oocysts in PBS. Suspensions are contained within a single capped bottle accessed by removing the top foil cap and bung.

### READ DIRECTIONS CAREFULLY

#### DIRECTIONS FOR USE

##### Contraindications:

- This product is contraindicated for vaccinating unhealthy birds.

##### Precautions:

- Avoid using in-feed or water anticoccidial medications at or soon after the time of vaccination, unless controlling clinical coccidiosis.
- Avoid vaccinating with EIMERIAVAX 3m unless birds are protected from exposure to litter from the previous batch for up to 14 days post-vaccination.
- Avoid using EIMERIAVAX 3m as a diluent for other medications or treatments.
- Avoid using in birds in lay.
- Avoid administration of sulfa drugs immediately following vaccination with EIMERIAVAX 3m as this will delay oocyst output and impair immune response.

##### Side Effects:

- Depression and some lesions associated with vaccination may be observed.

##### Dosage and Administration:

USE ALL PRODUCT WITHIN 24 HOURS OF OPENING. Birds from day-old can be vaccinated with EIMERIAVAX 3m. It is recommended that all birds be vaccinated prior to 18 weeks of age. The most appropriate age to vaccinate may be determined from knowledge of the challenge conditions applicable to the flocks concerned. The effect of EIMERIAVAX 3m vaccination during lay has not been assessed.

Eyedrop and spray method of administration is recommended for 1000 dose presentation only.

##### Eyedrop administration for the 1000 doses:

- Prior to use gently shake vials to fully resuspend all oocysts.
- Remove the cap and insert an eye-dropper tip or other administration device, calibrated so as to distribute

25 microlitre drops of vaccine. Hold each bird with its head tilted to one side.

- If using a dropper tip attached to the bottle, invert the bottle.
- Allow a single drop to form at the tip of the dropper or other application device and fall freely into the open eye, gently flooding it. The drop (before release) and dropper tip or administration device should NOT touch the eye surface.
- Allow the bird to blink before releasing it.
- Administer a second drop if the bird did not receive a full dose.

##### Spray method of administration for the 1000 doses:

EIMERIAVAX 3m has been demonstrated effective when administered using a hatchery spray apparatus that delivers 24 mL of diluted vaccine per box of 100 day-old chicks, i.e. 0.24 mL per chick. Please note the following points:

- The vaccine is to be diluted in water. If using a spray apparatus that delivers 24 mL per 100-chick box, one bottle of EIMERIAVAX 3m is diluted to a total volume of 240 mL.
- Open the vaccine bottle when required by peeling back the foil cap and removing the rubber bung. Rinse the vaccine bottle several times in the water diluent to ensure all the oocysts are removed from the bottle.
- Ensure the diluted vaccine is regularly agitated during the spray applications to ascertain that vaccine oocysts do not settle.
- The spray unit should deliver an even mist with the added blue dye (if used) evident. Ensure mist covers all chicks.
- After spraying, leave the chicks in a light place to allow preening and vaccine ingestion.
- Please contact Bioproperties if further information regarding specific spray units is required.

##### Spray method of administration for the 5000 doses:

EIMERIAVAX 3m has been demonstrated effective when administered using a hatchery spray apparatus that delivers 24 mL of diluted vaccine per box of 100 day-old chicks, i.e. 0.24 mL per chick. Please note the following points:

- The vaccine is to be diluted in water. If using a spray apparatus that delivers 24 mL per 100-chick box, one bottle of 5000 dose EIMERIAVAX 3m is diluted to a total volume of 1200 mL.
- Open the vaccine bottle when required by peeling back the foil cap and removing the rubber bung. Rinse the vaccine bottle several times in the water diluent to ensure all the oocysts are removed from the bottle.
- Ensure the diluted vaccine is regularly agitated during the spray applications to ascertain that vaccine oocysts do not settle.
- The spray unit should deliver an even mist with the added blue dye (if used) evident. Ensure mist covers all chicks.
- After spraying, leave the chicks in a light place to allow preening and vaccine ingestion.
- Please contact Bioproperties if further information regarding specific spray units is required.

##### Post vaccination general instructions:

- Continued exposure to oocysts shed in faeces is essential for the development of an effective immune response.
- Birds can be vaccinated from day-old. Anticoccidial medication is never recommended unless clinical coccidiosis from wild field strains is affecting bird performance. In these instances, manufacturer's recommendations for the medication program are to be followed.
- If depression associated with vaccination is noted, light intensity should be increased and/or extended and extra feed provided.
- Monitor birds carefully for 4 weeks post-vaccination. Some lesions can be expected. Lesions can be associated with an immune response and should not be used as the sole criterion for assessing efficacy.
- Investigate all suspect coccidiosis lesions as this vaccine does not protect against *Eimeria* species that are not represented in the vaccine. Monitor gross lesions scores for 4 weeks post-vaccination.

**Duration of Immunity:** An effective immune response can be expected for at least 18 weeks post-vaccination. Continuing exposure to oocysts in the environment is essential and provides ongoing stimulus to the immune system. Under these conditions, a protective immune response beyond 18 weeks of age can be expected. When birds are removed from the source of vaccine oocysts, such as when they are transferred between facilities, immunity may wane due to reduced exposure to cycling oocysts. As a result, birds may subsequently become susceptible to field infections from *Eimeria* species represented in the vaccine.

**WITHHOLDING PERIODS: Zero (0) days**

**USER SAFETY INFORMATION:**

Accidental contact of EIMERIAVAX 3m with eyes or mouth may cause irritation. Eyes should be rinsed immediately, and medical advice sought. The mouth should be rinsed two or three times with a mouth gargle.

Additional information is available on the product Safety Data Sheet.

**FIRST AID:**

First aid is not generally required. If in doubt, contact a Doctor or Poisons Information Centre on 13 11 26.

**DISPOSAL:**

Discard unused vaccine and empty vaccine bottles into a disinfectant solution (e.g. chlorine based bleach). Dispose of any such inactivated, unused vaccine and empty containers by wrapping in paper and putting in garbage.

In cases of spillage, soak up the liquid with an absorbent sponge or cloth and incinerate. Treat the surface with a disinfectant solution (e.g. chlorine based bleach).

**CAUTION:**

The capability of this vaccine to produce satisfactory results depends upon many factors including, but not limited to, conditions of storage and handling by the user, administration of the vaccine, health and responsiveness of individual birds and degree of field exposure. Therefore, directions for use should be followed carefully.

**CONTACT DETAILS:**

**BIOPROPERTIES Pty Ltd**  
**ACN 007 303 728**  
**36 Charter Street**  
**Ringwood**  
**Victoria 3134, Australia**  
**PH: 03 9871 2000 FAX: 03 9876 0556**

**BIOPROPERTIES Pty Ltd**  
**Eimeria Pty Ltd**



**STORAGE INSTRUCTIONS:**

Product should be stored between 2°C and 8°C (Refrigerate. Do not freeze).

**WARRANTY AND DISCLAIMER**

BIOPROPERTIES Pty Ltd manufactures, markets and sells EIMERIAVAX 3m on behalf of Eimeria Pty Ltd.

EIMERIAVAX 3m has been tested and meets safety and potency standards according to the Standards required by APVMA. BIOPROPERTIES Pty Ltd and Eimeria Pty Ltd accept no responsibility for events arising from the misuse or mishandling of

the product. BIOPROPERTIES Pty Ltd and Eimeria Pty Ltd give no warranty (express or implied) with respect to the product, including without limitation any warranty as to completeness, merchantability or fitness for a particular purpose. Under no circumstances shall BIOPROPERTIES Pty Ltd or Eimeria Pty Ltd be liable for indirect, special, consequential or punitive damages.

**APVMA No.** 63631/121757

**Code/Version No:** LFT- E3M/1,5 V2